

INDEX TO VOLUME XIV *

New names and the final members of new combinations are in **bold face type**

- Abies, 177, 181, 182; grandis, 174, 179, 180, 182, 183; pectinata, 8, 10
Absidia glauca, 149; septata, 149
Acer, 5, 6, 9, 181; circinnatum, 175, 198; macrophyllum, 174, 182, 183, 192; spicatum, 238
Achillea lanulosa, 118; Millefolium, 118
Achlya, 152, 154, 155, 166; prolifera, 146
Addition to the distribution of a rare fungus, 49
Aecidium, 105, 114; arcularium, 106; Pourthiacae, 293, 295
Agarics, Dark-spored — I, 61; — II, 121; — III, 200; — IV, 258
Agaricus, 27, 75, 125, 136, 200, 220, 269; abruptibulbus, 209; abruptus, 209; Achimenes, 220; acuminatus, 128; adnatus, 133; aeruginosus, 128; **alabamensis**, 202; amygdalinus, 220, 221; appendiculatus, 64; approximans, 215; argenteus, 211; Artemisiae, 75; arvensis, 208, 210; atomatus, 275; atrorufus, 263; brunescens, 211; bulbosus, 209; bulbularis, 131; bullaceus, 263; camarophyllus atramentosus, 48; camarophyllus fuliginosus, 48; campester, 136, 203, 205, 212; campester hortensis, 212; campestris, 136; campestris sylvicola, 209; Candolleanus, 71; capnoides, 74; chlamydopus, 205; Comaropsis, 75; **comtuliformis**, 203; comtulus, 202, 203; coprophilus, 259; cornilla, 131; coronatus, 71; cothurnatus, 205; cretaceus, 208, 221; cretaceus, 220; depilatus, 134; diminutivus, 189, 214; disseminatus, 270; distans, 126; echinatus, 213; elixus, 48; eludens, 217; eximius, 97; fabaceus, 220, 221; falcifolius, 278; fascicularis, 73; Feildeni, 141; flavescens, 27, 221; floridanus, 207; foederatus, 221; glutinosus, 122, 133; graciloides, 277; haemorrhoidarius, 215, 216, 219; halophilus, 212; hiascens, 278; hirtosquamulosus, 75; Hornemanni, 135; Howeanus, 141; hymenoccephalus, 64; hypoxanthus, 66; incertus, 64; Johnsonianus, 141; lacrymabundus, 67; lateritius, 72, 73; maculatus, 124; madeodiscus, 62; magniceps, 213; magnificus, 213; maritimus, 212; melanospermus, 131; melaspermus, 130, 131; merdarius, 138; micromegethus, 210; minutulus, 270; nitidipes, 75; obturatus, 141; odoratus, 276; olivaeosporus, 97; ornellus, 75; oxyosmus, 214; perplexis, 72; pholidotus, 268; pilosporus, 204; placomyces, 217, 218; polytrichophilus, 261; populinus, 66; praeimosus, 204; pratensis, 208; pusillus, 210; Rodmani, 207; **rubribrunescens**, 216; rupicola, 278; rutilescens, 211; saccharinophilus, 64; semiglobatus, 132; semivestitus, 269; solidipes, 203; sphaerosporus, 204; squamosus, 126; stercorarius, 133; Sterlingii, 217; Storcia, 66; subcernuus, 126; sublateritius, 72; subrufescens, 189, 206, 210, 219, 221; sylvaticus, 218; sylvicola, 209, 221; tabularis, 204; thraustus aurantiacus, 127; tricholepis, 76; umbonatescens, 128; variabilis, 206; velutinus, 67; violaceolamellatus, 71; viridulus, 128; viscidus, 122; xylogenus, 221
Agropyron divergens, 279; occidentale, 279; repens, 279; Smithii, 279; spicatum, 279

* It has been considered unnecessary to include here the species listed in the three following articles, since they are already indexed or specially listed.

Arthur: Uredinales collected by Fred J. Seaver in Trinidad. See p. 23, 24.

Murrill: Index to illustrations of fungi, XXIII-XXXIII. See p. 332.

Wann and Muenscher: A preliminary list of the myxomycetes of the Cayuga Lake Basin. See p. 39.

- Agrostis alba*, 279; *palustris*, 279
Aira caespitosa, 33, 34
Aleuria aurantia, 175
Aleurodiscus, 179; *amorphus*, 179;
subcruentatus, 179
Allescheria, 242; **Boydii**, 242
Alnus, 2, 5, 174, 175
Alopecurus aristulatus, 229, 230
Alternaria, 337
Amanita, 185; *calyptrata*, 185; *chlori-*
nosma, 185; *muscaria*, 185, 315;
pantherina, 92; *pantherinoides*,
185; *phalloides*, 317; *solitaria*, 185;
spissa, 92
Amanitopsis plumbea, 315; *vaginata*,
185; *vaginata alba*, 186; *vaginata*
livida, 186
Amauroderma, 42; *Gusmanianum*, 43;
Mosselmanii, 43; *picipes*, 43
Amelanchier, 2, 102, 289; *asiatica*,
288, 294; *Cusickii*, 91
American porias — I, Diagnoses of, 1
Amphisphaeria diffusa, 237; *subicu-*
losa, 237
Amsonia ciliata, 252, 297
Anellaria, 130; *fimiputris*, 130
Anemone, 176
Anisocarpua, 115
Another green-spored genus of gill-
fungi, 96
Aplospora, 336; *Nyssae*, 336
Aposphaeria brunneotincta, 102
Arachnion album, 195
Arcangeliiella caudata, 196
Aristida, 230
Armillaria, 187; *albolanaripes*, 186;
alphitophylla, 338; *corticata*, 187;
nardosmia, 25
Arnica, 114; *cana*, 114; *foliosa*, 114,
115
Artemisia, 118; *tridentata*, 45
Arthur, J. C., Uredinales collected by
Fred J. Seaver in Trinidad, 12
Articles, Notes and brief, 42, 90, 231,
335
Arundinaria Simoni, 86
Ascomycete isolated from a granular
mycetoma of man. Life history of
an undescribed, 239
Ascomycetes — I, Studies in tropical,
235
Ascophora mucedo, 143
Aster, 302, 306–310; *acuminatus*, 303,
309; *Chapmanii*, 304; *concinus*,
309; *conspicuus*, 254, 297, 303–306;
cordifolius, 246, 254, 303–306; *cor-*
rigiatus, 309; *divaricatus*, 303, 306;
dumosus, 303, 304; *ericoides*,
254, 303; *Fremontii*, 309; *fron-*
dosus, 309; *Geyeri*, 303–306;
hesperius, 254, 303; *hirsuti-*
caulis, 309; *juncus*, 309; *laevigatus*,
303; *laevis*, 246, 254, 303–306; *lae-*
vis Geyeri, 254; *lentus*, 254, 303,
304; *longifolius*, 297, 306; *Lowrie-*
anus, 303, 309; *macrophyllus*, 303–
306; *novi-belgii*, 304; *oblongifolius*,
309; *paniculatus*, 254, 303, 304, 306;
patens, 303; *patulus*, 309; *Pringlei*,
306, 309; *puniceus*, 303; *salicifolius*,
303; *schistosus*, 309; *spectabilis*,
309; *tenuicaulis*, 309; *undulatus*,
246, 254, 303–306; *vimineus*, 303,
304, 306; *viscosum*, 309
Atylospora, 258, 262, 264, 273, 278;
australis, 266, 278; *corrugis*, 267,
268; *microsperma*, 264; *multipe-*
data, 265; *prunuliformis*, 266, 278;
striatula, 267, 278; *umbonata*, 267,
268; *vestita*, 265
Auricularia nigrescens, 44
Bacidia inundata, 95
Bacterium marginatum, 336
Balsamorhiza, 112; *macrophylla*, 112
Barroetia, 107
Battarrea, 45; *laciniata*, 45
Betula, 5
Boehmeria nivea, 88
Boletinus cavipes, 181
Boletus, 45; *chrysenteron*, 180; *luri-*
us, 180; *luteus*, 25, 45, 180, 198;
scaber, 180; *scaber fuliginosus*, 180;
scaber testaceus, 180; *Zelleri*, 181
Borrchia arborescens, 111; *frutes-*
cens, 111
Botrydium, 95
Botrydiopodia Theobromae, 91
Bovista, 321; *montana*, 193; *plumbea*,
193
Bremia, 88
Brickellia, 107
Bromus, 279; *ciliatus*, 279
Broteria, 42
Bulgaria inquinans, 175
Bullaria, 105, 111, 112; *Kuhniae*, 106
Byssosphaeria diffusa, 237
Caeomus triannulatus, 111
Calamagrostis canadensis, 279
Calendula, 120; *officinalis*, 119
Calicium Curtisii, 103; **Rhois**, 103
Callistephus chinensis, 306
Calocera cornea, 178
Calonyction, 299; *aculeatum*, 297–
299; *grandiflorum*, 298
Calosphaeria, 100
Calvatia gigantea, 193; *lilacina*, 193;
lilacina occidentalis, 193; *rubro-*
flava, 193

- Calycadenia, 115
 Camarophyllus, 48
 Camillea Leprieurii, 338
 Campanula rapunculoides, 305
 Campanularius, 130
 Cantharellus aurantiacus, 188; cibarius, 317; floccosus, 188; infundibuliformis, 188
 Carduaceae, New or noteworthy rusts on, 104
 Carex, 104, 177; pendula, 33
 Carpinus, 1
 Carya illinoensis, 77
 Castanea dentata, 102
 Catastoma circumscissum, 193
 Cayuga Lake Basin, Myxomycetes of the, 38
 Cenangium balsameum, 101; **querneum**, 101; turgidum, 101
 Centromadia, 115
 Cephalosporium, 242; *Boydii*, 242
 Cercospora Brunkii, 91; fusimaculans, 198; minima, 89; piricola, 88; tortipes, 337; tuberculella, 337
 Cerotetium Dientrae, 336
 Chamaecyparis, 285; pisifera, 294; pisifera plumosa, 294; pisifera squarrosa, 294
 Chamonixia, 196, 197; caespitosa, 197
 Chanterel, 96; floccosus, 25, 43
 Characters of the southern tuckahoe, 222
 Chloroneuron, 96
 Chlorophyllum, 96
Chlorosperma, 96; **olivaespora**, 97
 Chrysopsis Mariana, 246, 248, 298, 303-306
 Cinnamomum camphora, 83
 Circinella, 166; conica, 149, 160-162; minor, 161-163
 Cladochytium, 84; Maydis, 84
 Cladoderris dendritica, 338
 Cladoniae, 95
 Cladosporium astericola, 337; herbarum, 240
 Clarkeinda, 61
 Clathrus locellus, 316
 Clavaria, 178; abietina, 178; botrytis, 178; corniculata, 178; cristata, 178; mucida, 178; subcaespitosa, 231; vermicularis, 178
 Cleavage in the sporangia of certain fungi, Method of, 143
 Clitocybe illudens, 43; odora, 187; phylophila, 26; subhirta, 26
 Coleosanthis, 106, 107; grandiflorus, 105
 Coleosporium, 114, 244-254, 297, 299, 302, 308; arnicale, 113-115; carneum, 250, 252, 308; delicatulum, 254-257; elephantopodis, 252; Helianthi, 244-253; inconspicuum, 244, 245, 247-253; Ipomoeae, 252, 297, 299-301; Laciniariae, 253; Madae, 114, 115; minutum, 252; ribicola, 299-302; Solidaginis, 250, 252, 256, 302, 303, 306-310; Terebinthinaceae, 250-253
 Coleosporium — I, Notes on some species of, 244; — II, 297
 Colletotrichum Boehmeriae, 88
 Collybia butyracea, 93; cirrata, 93; clusilis, 93; conigena, 93; distorta, 93; dryophila, 93; Hariolorum, 93; lilacina, 93; nummularia, 93; semitalis, 93; strictipes, 93; subdryophila, 93; tuberosa, 93; velutipes, 44
 Coniophora cerebella, 179
 Contributions to our knowledge of Oregon fungi — I, 173
 Convolvulus, 299; arvensis, 298; repens, 298; sepium, 298, 305
 Cookeina Tricholoma, 338
 Coprinus, 61, 220, 268, 271, 317, 318; atramentarius, 135; comatus, 135, 136; micaceus, 44; Spraguei, 274
 Coreopsis, 109, 110, 244, 245, 247, 248, 250, 251; delphinifolia, 251; lanceolata, 248, 251; major, 246-248, 250, 251, 298, 304; major Oemleri, 251; major rigida, 251; mexicana, 109, 110; mutica, 110; tinctoria, 303; tripteris, 246, 248, 251; verticillata, 246-248, 250-252, 298, 303-305
 Coriolopsis Copelandii, 93
 Coriolus sericeohirsutus, 44
 Cornus Nuttallii, 175
 Corticium lactescens, 178; salmonicolor, 91
 Cortinarius, 44; semisanguineus, 45
 Corylus californica, 174, 183
 Costantinella, 103; cristata, 103
 Craterellus cornucopioides, 178
 Crucibulum vulgare, 198
 Cryptoporus volvatus, 181
 Cryptosporella anomala, 46
 Cultures of heteroecious rusts, 1920-1921, 228
 Cyathus stercoreus, 198; vernicosus, 198
 Cydonia japonica, 285, 294; sinensis, 285; vulgaris, 282, 284, 285, 294
 Cyllindrophora, 337
 Cyllindrosporium acerinum, 198
 Cyphella Heveae, 91; marginata, 179
 Cystopus, 146
 Dacryomyces aurantia, 177

- Daldinia vernaica*, 174
 Dark-spored agarics — I, 61; — II, 121; — III, 200; — IV, 258
Dasyscypha conferta, 118; *Millefolii*, 118; *Synedrella*, 117
 Davis, W. H., *Urocystis Agropyri* on redtop, 279
Deconica, 258, 268; *atrurufa*, 263; *bryophila*, 261; *bulbosa*, 260; *bulbacea*, 258, 262, 263; *coprophila*, 259; *nuciseda*, 259; ***polytrichophila***, 261; ***pyrispora***, 261; *rhomboidospora*, 259; *scatigena*, 258; *semistriata*, 260; *subviscida*, 262; ***tomentosa***, 259
Delitescor, 258
Dendrostilbella, 241, 242; *Boydii*, 242
Derbesia, 148
 Diagnoses of American porias — I, 1
Diaporthe Menispermii, 233; *Triostei*, 233
Diatrype, 174
Diatrypella, 100; ***ciliatula***, 100
Dicaeoma Balsamorhizae, 112; *Brickelliae*, 106; *cubense*, 116, 117; *Eriophorii*, 119; *Hemizoniae*, 115; *Kuhniae*, 106; *nudum*, 115; *Sarcobati*, 229; *Synedrella*, 116; *trianulatum*, 111
Didymella Agrostidis, 233
Diervilla Lonicera, 101
Dimorphotheca, 119, 120; *cuneata*, 119
Discina ancilis, 175
Distichlis stricta, 229, 230
 Distribution of a rare fungus, An addition to the, 49
Dodecatheon pauciflorum, 229, 230
Drosophila, 61, 121, 138, 139, 150, 264; *appendiculata*, 63, 64, 71, 141, 265; *atrofolia*, 71; ***delineata***, 68; ***echiniceps***, 68, 69; ***fragilis***, 64; ***hololanigera***, 70; *hydrophila*, 72; *lacrymabunda*, 67, 69; ***madeodisca***, 62, 65, 72; ***Peckiana***, 68; ***pecosense***, 63; ***rigidipes***, 70; *Storea*, 66, 72
Echinodontium tinctorum, 180
Electra, 110; *Galeottii*, 110
Elephantopus, 250; *carolinianus*, 248–250, 298, 303, 304; *nudatus*, 298; *tomentosus*, 249, 254, 298, 303–306
Eleutheranthera ruderalis, 116, 117
 Elliott, J. A., Some characters of the southern tuckahoe, 222
Elymus, 279; *arenarius*, 279; *canadensis*, 279; *robustus*, 279; *virginicus*, 279
Emilia sagittata, 119; *sonchifolia*, 119
Equisetum, 175
Eriophorum, 119
Eupatorium, 107, 108; *atriplicifolium*, 107; *maculatum*, 254; *pansamalense*, 108; *urticaefolium*, 108
Eurotiopsis, 242; *Gayoni*, 241
Eurotium, 241
Euthamia, 254–256; *caroliniana*, 255–257; *graminifolia*, 254–257, 306; *leptophylla*, 257
Excidia recisa, 177; *Zelleri*, 177
Fagus, 1, 2, 5
 Farlowianae, Reliquiae, 99
 Fink, B., An addition to the distribution of a rare fungus, 49; A new lichen from an unusual substratum, 95
 Fitzpatrick, H. M., Thomas, H. E., and Kirby, R. S., The *Ophiobolus* causing take-all of wheat, 30
Flaveria angustifolia, 118; *campestris*, 118
Fomes annosus, 181; *applanatus*, 181; *connatus*, 9; *igniarius*, 94, 181; *lamaensis*, 93; *leucophaeus*, 181; *lignosus*, 91; *lineatus*, 92; *pinicola*, 181; *populinus*, 9; *pseudoferreus*, 91; *roseo-albus*, 92, 93; *roseus*, 182, 183; *semilaccatus*, 92
 Fragrant polypore, A, 46
 Fraser, W. P., Cultures of heteroeccious rusts, 1920–1921, 228
 Fungi — I, Contributions to our knowledge of Oregon, 173
 Fungi, Illustrations of, 25
 Fungi, Index to illustrations of, 332
 Fungi, New Japanese, 81, 282
 Fungus, An addition to the distribution of a rare, 49
Fusarium, 337; *oxysporum*, 51
Galerula Hypni, 27
Ganoderma, 92; *Curranii*, 92; *leptopum*, 92; *oregonense*, 43, 92; *umbraeculum*, 92
Gaultheria Shallon, 178
Gautieria, 196; *morchelliformis*, 196; ***Parksiana***, 196, 197
Geaster, 311, 316, 321; *delicatus*, 191; *hygrometricus*, 192; *hygrometricus giganteus*, 192; *lagenaeformis*, 192; *Morganii*, 192; *saccatus*, 192
Gelatinosporium abietinum, 101
 Genus of gill-fungi, Another green-spored, 96
Geophila, 125; *capnoides*, 74
Geopyxis cupularis, 175
 Gill-fungi, Another green-spored genus of, 96

- Glaux, 229; maritima, 228-230; Michauxiana, 228
 Gleditsia, 2
 Gliocladium penicillioideus, 101
 Gloeosporium acutilobum, 233; Balsameae, 337; bicolor, 337; limetticola, 335
 Glomerella, 84; Cinnamomi, 82
 Gnomonia Coryli, 174
 Gnomoniopsis acerophila, 233
 Godronia Nemopanthidis, 101; **turbinata**, 101
 Gomphidius, 121, 200; flavipes, 124; furcatus, 125; glutinosus, 44, 122; jamaicensis, 121; maculatus, 124, 125; nigricans, 123; oregonensis, 121, 191; tomentosus, 121; vinicolor, 121, 123; viscidus, 44, 122, 124
 Green-spored genus of gill-fungi, Another, 96
 Grossularia, 301; Cynosbati, 301; divaricata, 300, 302; hirtella, 300, 302; inermis, 300-302; innominata, 300, 302; leptantha, 300, 301; missouriensis, 300, 302; reclinata, 300-302; setosa, 301
 Guignardia Cinnamomi, 84
 Gymnochilus roseolus, 268
 Gymnomyces Gardneri, 198; pallidus, 197, 198
 Gymnopilus flavidellus, 28; polychrous, 75, 141, 142
 Gymnopus velutipes, 45
 Gymnosporangium, 283-286, 290, 291, 293; alpinum, 294; amelanchieris, 289; asiaticum, 282-285, 287, 292, 294; blasdaleanum, 295; chinensis, 287, 293, 294; clavariaeforme, 295; clavariiforme, 287, 289, 293; clavariaeforme, 287, 293; clavipes, 289; confusum, 284, 293, 294; Haraeae, 283, 284, 287, 290, 294; hemisphaericum, 289, 294; Idetae, 287, 294; japonicum, 283-285, 288, 293, 294; Juniperi, 294; juniperinum, 289, 294; koreaense, 294; koreaensis, 284, 287; Miyabei, 288, 294; Photinae, 284, 285, 293; Sabiniae, 295; Shiraianum, 291, 294; spiniferum, 284, 294; tremelloides, 288, 294; Yamadae, 285, 287, 293; Yamadae, 286, 293
 Gyrocephalus rufus, 177
 Gyromitra esculenta, 176
 Harpaecarpus, 115
 Hebeloma, 75; luteum, 29
 Hedgcock, G. G., and Hunt, N. R.,
 Notes on some species of Coleosporium — I, 244; — II, 297
 Helianthus, 244, 245, 247, 250; angustifolius, 298, 304; annuus, 255, 304; australis, 250; decapetalus, 246-248, 250, 251, 298, 305; divaricatus, 246-248, 250, 251, 254, 298, 304, 305; doroconoides, 250; Dowellianus, 246, 248; Eggertii, 250; giganteus, 246, 247, 250; glaucus, 246-248, 250, 251; grosseserratus, 250; hirsutus, 246-248, 250; microcephalus, 246-248, 250, 251; occidentalis, 248, 250, 254, 303, 304; Radula, 246, 247, 251, 298; saxicola, 251; strumosus, 251; tuberosus, 248, 251, 304
 Helminthosporium macrocarpum, 82; Oryzae, 81, 82
 Helvella crispa, 176, 198; elastica, 176; infula, 176; lacunosa, 176
 Hemizonella, 114, 115
 Hemizonia, 114, 115
 Herpotrichia, 236, 237; albidostoma, 235, 236, 238; diffusa, 236-238; incisa, 238; rhodomphala, 237; rhodospiloides, 236, 237
 Heteroecious rusts, Cultures of, 228
 Heteromeles, 196
 Hevea brasiliensis, 91
 Hexagonia carbonaria, 183
 History of mycological illustration, A sketch of the, 311
 Hunt, N. R., and Hedgcock, G. G.,
 Notes on some species of Coleosporium — I, 244; — II, 297
 Hydnangium carneum, 335
 Hydnotrya ellipsospora, 176
 Hydnum auriscalpium, 180; erinaecus, 180; ochraceum, 180; repandum, 180; subfuscum, 180
 Hydrocybe ceracea, 44; Earlei, 44
 Hydrodictyon, 149
 Hygrophorus, 48; caprinus, 48; eburneus, 188; fuliginosus, 48; miniatus, 188; miniatus sphagnophilus, 188
 Hylocomium triquetrum, 188
 Hymenochaete spreta, 179
 Hymenopappus carolinensis, 118
 Hypholoma, 61, 72, 75, 96, 121, 190; aggregatum, 66; aggregatum sericeum, 66; appendiculatum, 63, 64, 71; Artemisiae, 75; Boughtoni, 67, 68; Candolleum, 63, 71; capnoides, 72, 74, 191; comaropsis, 75; comatum, 71; confertissimum, 71; coronatum, 71; cutifractum, 64; delineatum, 68; echiniceps, 69; fasciculare, 72-74, 190, 191; flocculentum, 64; fragile, 64; hololanigerum, 70;

- hydrophilum, 191; lacrymabundum, 66, 67, 191; lateritium, 72; madeo-discum, 62, 63, 71; nitidipes, 75; ornellum, 75; Peckianum, 68; pe-cosense, 63, 133; perplexum, 73; populinum, 72; Pseudostorea, 66; rigidipes, 70; rugocephalum, 67, 68; subaquilum, 62, 63, 71; sublateri-tium, 73; velutinum, 191; veluti-num leiocephalum, 71; vinosum, 97
 Hypoxylon atropunctatum, 174; prui-natum, 51
 Illustration, A sketch of the history of mycological, 311
 Illustrations of fungi, XXXIII, 25
 Illustrations of fungi, Index to, 332
 Index to American mycological liter-ature, 53
 Index to illustrations of fungi, XXIII-XXXIII, 332
 Inocybe, 91, 214, 232, 260; fastigiata, 189
 Inonotus dryophilus, 44; perplexus, 43
 Ipomoea, 299; Batatas, 298; carolini-ana, 298, 299; fistulosa, 303; lacu-nosa, 298, 299; pandurata, 298, 299, 303; triloba, 298, 299
 Iris foetidissima, 33-35
 Irpex, 1; lacteus, 180
 Jackson, H. S., New or noteworthy rusts on Carduaceae, 104
 Japanese fungi, New, 81, 282
 Juniperus chinensis, 282, 283, 285, 286, 290, 293, 294; chinensis pro-cumbens, 282, 286, 293, 294; littor-alis, 292-294; nana, 294, 295; nip-ponica, 294; rigida, 285, 288, 293, 294
 Justicia procumbens, 87
 Kirby, R. S., Fitzpatrick, H. M., and Thomas, H. E., The Ophiobolus causing take-all of wheat, 30
 Knowledge of Oregon fungi—I, Con-tributions to our, 173
 Krieger, L. C. C., A sketch of the history of mycological illustration (higher fungi), 311; Schizophyllum commune with a stipe, 47
 Kuhn, 107
 Laccaria laccata, 44
 Laciniaria acidota, 303; elegans, 246, 248, 298; elegantula, 298; gracilis, 298; graminifolia, 252, 298; pyc-nostachya, 298; tenuifolia, 298
 Lactarius pyrogalus, 188
 Laetiporus sulphureus, 44
 Lagenidium entophyllum, 145
 Lagophylla, 115
 Lapsona communis, 176
 Lasiosphaeria strigosa, 174
 Lentinula detonsa, 338
 Lentinus, 223
 Lentodiellum concavum, 338
 Lenzites saepiaria, 93, 188; vialis, 189
 Lepiota, 96, 186, 214, 215; amian-thina, 214; clypeolaria, 186; cre-tacea, 44; granulosa, 186; Mor-gani, 96; naucina, 221; procera, 43, 315; **pulcherrima**, 186; rubro-tincta, 186; rubrotinctoides, 186
 Leptosphaeria Collinsoniae, 233
 Leptothyrium conspicuum, 233
 Lichen from an unusual substratum, A new, 95
 Licipenicillium insigne, 101
 Life history of an undescribed asco-mycete isolated from a granular mycetoma of man, 239
 Lilliputia Gaillardii, 101
 List of the myxomycetes of the Ca-yuga Lake Basin, 38
 Lophodermium Thuyae, 337
 Lycoperdon, 321; cruciatum, 192; ele-gans, 192; fuscum, 192; gemmatum, 192; pratense, 192; pusillum, 192; pyriforme, 192; rimulatum, 192; solidum, 320; Wrightii, 193
 Macbride's North American slime-moulds, 233
 Macrosporium, 44, 337
 Madaria, 115
 Madia, 114, 115; Nuttallii, 115
 Marasmius plicatulus, 188
 Meeting, The Toronto, 50
 Melampodium divaricatum, 115, 117
 Melanconium oblongum, 46
 Melanoleuca eduriformis, 27; suban-nulata, 187; Thompsoniana, 26
 Melanotus, 61
 Meliola Abietis, 174
 Merulius brassicaefolius, 184; **pilo-sus**, 184, 185; rugulosus, 185; tremulosus, 185
 Method of cleavage in the sporangia of certain fungi, 143
 Microdiplodia Populi, 232
 Micropuccinia, 108, 109, 118, 119; conferta, 119; Millefolii, 119; Synedrellae, 117
 Miles, L. E., A new species of Myri-angium on pecan, 77
 Monascus purpureus, 42

- Monocytrium, 148
 Morchella, 316; angusticeps, 175; crassipes, 175, 316; deliciosa, 175; esculenta, 315, 316; semilibera, 176
 Mucor, 143, 144; mucedo, 144, 146, 163, 164, 166; racemosus, 161; spinescens, 149, 161
 Muenscher, W. C., and Wann, F. B., A preliminary list of the myxomycetes of the Cayuga Lake Basin, 38
 Murrill, W. A., A fragrant polypore, 46; Another green-spored genus of gill-fungi, 96; Dark-spored agarics — I, 61; — II, 121; — III, 200; — IV, 258; Hygrophorus caprinus, 48; Illustrations of fungi — XXXIII, 25; Index to illustrations of fungi, XXIII-XXXIII, 332; The Toronto meeting, 50
 Mycena palustris, 93; pura, 44
 Mycetoma of man, Ascomycete isolated from a granular, 239
 Mycogone puccinioides, 198; roseola, 198
 Mycological illustration, A sketch of the history of, 311
 Mycosphaerella, 232; Arundinarieae, 86; bambusifolia, 85
 Myriangium, 80; **tuberculans**, 80
 Myriangium on pecan, New species of, 77
 Myxomycetes of the Cayuga Lake Basin, 38
 Naematelia encephala, 177
 Naucoria, 75; pennsylvanica, 75; semiorbicularis, 189
 Nectria, 174; coccinea, 174, 175; Coryli, 174; episphaeria, 174; galligena, 174
 Neopecticia, 236, 237; diffusa, 235-237
 Neurolaena lobata, 119
 Neurophyllum viride, 96
 New Japanese fungi, Notes and translations — XI, 81; — XII, 282
 New lichen from an unusual substratum, 95
 New or noteworthy rusts on Carduaceae, 104
 New Septobasidium on Pinus Strobus, 55
 New species of Myriangium on pecan, 77
 North American slime-moulds, Macbride's, 233
 Notes and brief articles, 42, 90, 231, 335
 Notes on some species of Coleosporium — I, 244; — II, 297
 Noteworthy rusts on Carduaceae, 104
 Nyssa, 336
 Odontia fragilis, 180
 Olpidiopsis, 146, 147, 150-152; major, 151; Saprolegniae, 152, 154, 155
 Olpidium Dicksonii, 148, 151; Viciae, 152
 Oospora, 337
 Ophiobolus, 30-32, 35; cariceti, 35, 36; eucryptus, 32; graminis, 30-33, 35, 36; herpotrichus, 30, 36
 Ophiobolus causing take-all of wheat, 30
 Oregon fungi — I, Contributions to our knowledge of, 173
 Oryza sativa, 81
 Ostrya, 5
 Otidea leporina, 175
 Overholts, L. O., Diagnoses of American porias — I, 1
 Oxalis oreghana, 198
 Pachyma Cocos, 92, 222, 223
 Panaeolus, 130, 136; campanulatus, 275; epimyces, 135
 Panicum, 198; proliferum, 89; proliferum acuminatum, 89
 Parthenice mollis, 108
 Parthenium, 252; argentatum, 109; Hysterophorus, 109, 116; incanum, 109; integrifolium, 246, 252, 253, 303, 304
 Paxillus pubescens, 122
 Pecan, A new species of Myriangium on, 77
 Penicillium insigne, 101
 Peniophora cinerea, 179; glabra, 179; glebulosa, 179; incarnata, 179
 Peridermium, 302; acicolum, 302, 310; delicatulum, 254; inconspicuum, 244; montanum, 302, 310
 Pezicula **pruinosa**, 102
 Peziza, 102; Acetabulum, 317; quereana, 101
 Phacidium Balsameae, 337; expansum, 337; planum, 337
 Phaeosphaeria Bambusae, 86; Oryzae, 86
 Pharbitis, 299; barbigera, 298, 299; hederacea, 298, 299; purpurea, 298, 303, 304
 Phialea alniella, 175
 Pholiota, 42, 141, 214, 220; duroides, 75; Howeana, 141; Johnsoniana, 141; ornella, 75; spectabilis, 189
 Phomopsis Impatiensis, 233
 Photinia, 284; laevis, 293; villosa, 283, 284, 293, 295

- Phragmidium, 317
 Phycomyces, 149, 166; nitens, 149, 158, 166
 Phyllostachys bambusoides, 85; puberula, 85
 Phyllosticta, 86, 233
 Physoderma Maydis, 84; Zeae-Maydis, 85
 Phytophthora, 146, 336; Faberi, 91; infestans, 148, 336
 Piggottia Vaccinii, 337
 Pilobolus, 148, 149, 158, 159, 162, 166, 167; crystallinus, 148, 160, 161, 163-168
 Pilosace, 61, 72, 75, 76, 96, 121, 136; algeriensis, 76, 136; eximia, 76; Peckii, 97
 Pinus, 250-252, 256, 301; Banksiana, 250; Bungeana, 300; canariensis, 256; caribaea, 247, 255-257, 299, 300, 306, 307, 309; clausa, 255; contorta, 255-257, 302, 305, 306-309; Coulteri, 255-257, 306, 307, 309; densiflora, 255; edulis, 247, 255, 256, 300-302, 306; echinata, 248, 250-252, 255-257, 297, 299, 303, 306-309; Girardiana, 300; glabra, 247, 248, 255-257; heterophylla, 309; Lambertiana, 56; Mayriana, 255-257, 300, 306; monophylla, 255, 300; montana, 255, 306; monticola, 56, 58, 59; nigra, 307; nigra austriaca, 309; nigra poiretiana, 256; palustris, 248, 251, 252, 254-257, 298, 306, 307; Pinca, 300, 302; pungens, 304, 306-308; radiata, 247, 305, 307, 309; resinosa, 255, 256, 305, 307; rigida, 252, 254-257, 300, 303, 306-309; scopulorum, 255-257, 306, 308, 309; serotina, 252, 255-257, 300, 306; strobiformis, 300; Strobis, 58, 300; taeda, 252, 255-257, 298, 300, 304, 306-309; Thunbergii, 255; virginiana, 245-248, 250, 251, 300
 Pirus, 295; Aria kamaoensis, 294; Aucuparia, 293, 294; Aucuparia japonica, 294; communis, 1, 88, 285, 294; Malus, 286, 290, 293; Miyabei, 294; spectabilis, 286, 287, 293; sinensis, 88, 282, 284, 290, 291, 293, 294; Toringo, 285-287, 290, 293; Zumi, 289, 290, 294
 Plantago, 230; eriopoda, 230
 Plasmopara Wildemaniana macrospora, 87
 Pleomassaria Magnoliae, 100; maxima, 100
 Pleonectria berolinensis, 175
 Pleurotus, 187, 232; ostreatus, 44; sapidus, 188; serotinus, 188
 Pluteus, 76, 98
 Polyphagus Euglenae, 147
 Polypore, A fragrant, 46
 Polyporus anebus, 92; arcularius, 182; basilapiloides, 43; Bracei, 44; Broomei, 10; caliginosus, 92; carbonarius, 91; chioneus, 182; cutifRACTUS, 91; dryadeus, 182; fastuosus, 92; floriformis, 182; fragilis, 182; galactinus, 182; hirsutulus, 182; hirtus, 182; lucidus, 93; Macounii, 182; Mariannus, 92; minor-Myliittae, 43; Myliittae, 43; pargameus, 3, 4; picipes castaneus, 182; pubescens, 182; rhodopheus, 92; rigidus, 9, 11; rufescens, 183; sensibilis, 183; Spraguei, 183; squamosus 315; trabeus, 182; tulipiferus, 4; tumulosus, 43; versicolor, 182; volvatus, 94; zonatus, 183
 Polysaccum, 321; turgidum, 194
 Polythelis Thalictri, 177
 Polytrichum, 261
 Populus, 5, 181, 184; grandidentata, 43, 47
 Poria, 1, 183; ambigua, 1, 2, 4; Broomci, 11; contigua, 183; ferruginosa, 5-7; Macouni, 6; marginella, 7; nigrescens, 7, 11; odora, 10, 11; sanguinolenta, 183; tenuis, 6; undata, 10; viticola, 6, 7
 Porias, Diagnoses of American, — I, 1
 Poronidulus conchifer, 51
 Pourthiaea villosa, 284
 Pratella, 200
 Preliminary list of the myxomycetes of the Cayuga Lake Basin, 38
 Protococcus, 95
 Protomyces bellides, 147
 Prunulus, 266
 Prunus, 5; demissa, 91
 Psalliota, 200
 Psathura, 264
 Psathyra, 96, 264, 269, 278; australis, 278; gyroflexa, 274; microsperma, 264; multipedata, 265; obtusata, 268; persimplex, 268; pholidota, 268; polytrichophila, 261, 268; prunuliformis, 278; roseola, 268; semivestita, 265, 269; silvatica, 269; striatula, 278; umbonata, 267; vestita, 265, 269
 Psathyrella, 258, 264, 269, 271; angusticeps, 275; atomata, 272, 275; Bartholomaei, 272, 273; betulina, 273; castaneicolor, 272; Clementsii, 274; crenata, 278; debilis, 274, 278; disseminata, 270; distantifolia,

- 277; *falcifolia*, 278; *gracilis*, 278; *gracillima*, 274; *graciloides*, 276, 277; *hiascens*, 278; *hirta*, 272; *leucostigma*, 271; *minima*, 272; *minutula*, 269, 270; *odorata*, 276; ***petasiformis***, 276; *rupicola*, 278; *sulcata*, 274; *tenera*, 271
- Pseudodipidium* *Aphanomycis*, 148, 151, 154, 156, 158
- Pseudoplectania* *melaena*, 175
- Pseudospora* *Lindstedtii*, 148
- Pseudotsuga*, 176, 177, 181, 196; *taxifolia*, 43, 176, 178-181, 184, 186, 196
- Psilocybe*, 138, 258, 263, 264; *atomatoides*, 277; *conissans*, 72
- Puccinia*, 104-106, 108, 114, 115, 120; *abrupta*, 113; *Actinellae*, 118; *additicia*, 109, 110; *artemisiicola*, 118; *Asteris*, 118; *aut-ou-Puccinia*, 105; *Balsamorhizae*, 111, 112; *Barroetiae*, 106; *brachy-Puccinia*, 106, 111, 112; *Brickelliae*, 106; *Burnettii*, 176; ***clara***, 104; *cognata*, 113; *conferta*, 118; *Coreopsisidis*, 109, 110; *Diaziana*, 116, 117; *Distichlidis*, 228-230; *Electrae*, 110; *Eleutherantherae*, 116, 117; *Emiliae*, 116, 119, 120; *Eriophorii*, 119; *Eupatorii*, 107; *ferox*, 113; ***Flaveriae***, 117; *gigantispora*, 176; *Grindeliae*, 118; *Hemizoniae*, 114, 115; *hyalina*, 104; *invelata*, 113; *irregularis*, 113; *Kuhniae*, 106, 107; *Lagophyllae*, 115; *Lapsanae*, 176; *Madiae*, 114, 115; *Millefolii*, 118; *mirifica*, 111; *nuda*, 113-115; *pachyspora*, 107; ***Parthenices***, 108; *Parthenii*, 109; *praealta*, 105; *recondens*, 119; *recondita*, 118; ***redempta***, 107; *Silphii*, 116; *solida*, 116, 117; *spilanthicola*, 116; *subdecora*, 105; *subnitens*, 228-230; *Synedrellae*, 115, 116; *Tetranthi*, 116, 117; *tolimensis*, 108; *Tonduziana*, 104, 105; ***triannulata***, 111; *Tridacis*, 116, 117; *turgidipes*, 110; ***vaga***, 112, 113; *Verbesinae*, 113; *Wyethiae*, 111, 112; *Zinniae*, 116, 117
- Pucciniastrum*, 46; *americanum*, 46; *arcticum*, 46
- Pyropolyporus* *Williamsii*, 92
- Pythium* *entophyllum*, 145; *intermedium*, 156, 158; *proliferum*, 157
- Quamoclit* *coccinea*, 298, 299; *Quamoclit*, 298
- Quercus*, 2, 5, 176; *Garryana*, 174, 175, 183, 195, 196, 198, 199
- Radulum* *Owensii*, 180
- Ramularia* *cilinodis*, 337; *Eamesii*, 233; *minax*, 337; *Oxalidis*, 198
- Ranunculus* *apetalus*, 229, 230
- Redtop*, *Urocystis* *Agropyri* on, 279
- Reliquiae* *Farlowianae*, 99
- Rhaphidophora* *graminis*, 36
- Rhizopogon* *diplophloeus*, 195; *induratus*, 195; *maculatus*, 196, 335; *roseolus*, 196; *rubescens*, 196
- Rhizopus*, 143, 144, 337; *nigricans*, 143, 149, 158-160, 163-166, 168
- Rhodochytrium*, 147
- Ribes*, 175, 301; *alpinum*, 300; *americanum*, 300-302; *aureum*, 300-302; *coloradense*, 301; *fasciculatum*, 300, 302; *inebrians*, 300-302; *malvacum*, 300; *mescalerium*, 301; *montigenum*, 301; *nigrum*, 254, 300, 302; *odoratum*, 300-302; *pumilum*, 301; *sanguineum*, 301; *vulgare*, 300, 302; *Wolfii*, 301
- Robinia* *Pseudacacia*, 1
- Roestelia*, 282, 283; *cancellata*, 295; *koreaensis*, 282-284, 294; *Photinia*, 284, 293; *Pourthiaecae*, 293; *solenoides*, 294; *solitaria*, 294
- Roskovites* *granulatus*, 45
- Rozella*, 146, 147, 150, 151
- Rubus*, 190; *idacus* *aculeatissimus*, 46; *occidentalis*, 46; *pubescens*, 46; *strigosus*, 46; *triflorus*, 46
- Rudbeckia* *laciniata*, 246
- Russula*, 45; *emetica*, 45; *foetens*, 315; *nigricans*, 315; *ochrophylla*, 188; *virescens*, 315
- Rusts*, *Cultures of heteroecious*, 228
- Rusts on Carduaceae*, *New or noteworthy*, 104
- Saccardinula* *Alni*, 233
- Saccharum* *Narenga*, 87
- Salix*, 5, 174, 175; *alba*, 47
- Sambucus* *glauca*, 177; *pubens*, 277
- Sanicula* *Menziesii*, 174
- Saprolegnia*, 146-148, 150-152, 154, 155, 158-160, 163, 166
- Sarcoseypha* *coccinea*, 175
- Sasa* *paniculata*, 86
- Schizonella* *melanogrammia*, 177
- Schizophyllum*, 47, 232; *alneum*, 48; *commune*, 47, 232; *commune stipitatum*, 48
- Schizophyllum* *commune* with a stipe, 47
- Schulzeria*, 96
- Schwarze*, C. A., *The method of cleavage in the sporangia of certain fungi*, 143
- Scirpus* *maritimus*, 229

- Scleroderma, 194, 321; *aurantiacum*, 194; *cepa*, 193, 194; *hypogaeum*, 193, 194
- Scleroderma, 102; **Cephalanthi**, 102
- Sclerotinia, 91; *demissa*, 91; *gregaria*, 91
- Seaver, F. J., Macbride's North American slime-moulds, 233; Studies in tropical ascomycetes — I, 235
- Seaver, Fred J., Uredinales collected by, 12
- Secotium tenuipes, 335
- Senecio, 119, 120; *aureus*, 254, 303, 305; *Cineraria*, 120; *obovatus*, 303, 305
- Sepedonium chrysospermum, 198
- Septobasidium, 55-57; **pinicola**, 55-58; *pseudopedicellatum*, 56
- Septobasidium on Pinus Strobus, A new, 55
- Septocladia dichotoma, 231
- Septoria Acetosella, 233; *Coreopsisidis*, 337; *hieracicola*, 233
- Shear, C. L., Life history of an undescribed ascomycete isolated from a granular mycetoma of man, 239
- Sidalcea, 176
- Silphium, 116, 250, 252; *angustatum*, 253; *aspermum*, 253; *asteriscum*, 246, 250, 252, 253; *compositum*, 253; *dentatum*, 253; *glabrum*, 253; *gracile*, 253; *integrifolium*, 246, 248, 250, 252, 253; *laciniatum*, 253; *laevigatum*, 253; *perfoliatum*, 246; *pinnatifidum*, 253; *scaberrium*, 253; *Simpsonii*, 298; *terebinthinaceum*, 253, 298; *trifoliatum*, 246, 252, 253
- Sketch of the history of mycological illustration (higher fungi), 311
- Slime-moulds, Macbride's North American, 233
- Snell, W. H., A new Septobasidium on Pinus Strobus, 55
- Solidago, 250, 302-305, 307-309; *amplexicaulis*, 309; *austrina*, 309; *bicolor*, 249, 254, 303-307, 309; *Boottii*, 309; *brachyphylla*, 309; *caesia*, 303, 304; *canadensis*, 247, 254, 298, 303-306, 309; *celtidifolia*, 309; *Chandonnetii*, 309; *Chapmannii*, 303, 304, 306, 309; *concinna*, 309; *Curtissii*, 309; *decumbens*, 309; *Drummondii*, 309; *erecta*, 254, 303, 306, 309; *fistulosa*, 298, 303, 304, 306, 309; *glomerata*, 309; *hispida*, 303, 306, 309; *junceae*, 249, 254, 298, 303-306, 309; *lancifolia*, 309; *mon-*
- ticola*, 305, 309; *multiradiata*, 254, 298, 303-306, 309; *neglecta*, 304, 306, 309; *nemorialis*, 304; *odora*, 309; *petiolaris*, 309; *pinensis*, 309; *Pinetorum*, 309; *pulverentula*, 309; *pulverula*, 309; *Purshii*, 309; *Riddellii*, 303-306, 309; *rigida*, 304, 309; *rigidiuscula*, 309; *rugosa*, 254, 255, 298, 303, 304, 306, 309; *rupes-tris*, 309; *sempervirens*, 304, 306; *serotina*, 303, 304, 306, 309; *speciosa*, 254, 255, 303, 304, 306, 309; *squarrosa*, 303-306, 309; *stricta*, 309; *tortifolia*, 304, 309; *unigulata*, 309; *Vaseyii*, 309
- Some characters of the southern tuckahoe, 222
- Sorbus, 295; *Aucuparia*, 293; *japonica*, 293, 294; *sambucifolia pseudo-gracilis*, 294
- Sparassis radicata, 177
- Spartina Michauxiana, 228; *pectinata*, 228, 230
- Spathularia clavata, 176
- Species of Coleosporium — I, Notes on some, 244; — II, 297
- Species of Myriangium on pecan, A new, 77
- Sphaeria, 102; *albidostoma*, 238; *cariceti*, 32-36; *ciliatula*, 100; *diffusa*, 237; *eucrypta*, 32-35; *rhodomphala*, 237
- Sphaerobolus stellatus, 198
- Sphaeronema pruinosa, 102
- Sphaeropsis ulmicola, 335
- Sphaerulina pallens, 337
- Sphagnum cuspidatum, 103; *Torreyanum*, 103
- Spilanthes oleracea, 116
- Sporangia of certain fungi, Method of cleavage in the, 143
- Sporidesmium Fusus, 100
- Sporidinia, 146, 148, 149, 159; *grandis*, 148, 158-164, 166-168
- Stagonospora Meliloti, 233; *tetrame-ra*, 337
- Stammaria Persoonii, 175
- Stemphylium, 44
- Stereum fuscum, 178; *purpureum*, 178
- Stichospora Madiae, 115
- Stieronema ciliatum, 228
- Stilbum Rhois, 103
- Stropharia, 76, 121, 125, 136, 200; **acuminata**, 128; **adnata**, 133; *aeruginosa*, 128, 140, 142, 190; *albocyanea*, 140; *albonitens*, 140; *ambigua*, 190; **annellariiformis**, 129; *bilamellata*, 130, 131, 137, 211; *caesifolia*, 136, 137; *caesiospora*,

- 134; *campestris*, 136; *coprinophila*, 135; *coronilla*, 131, 137, 139, 141; *cotonea*, 66; *depilata*, 134; *distans*, 126; *elegans*, 140; *epimyces*, 76, 135; *Feildenii*, 141; *Hardii*, 134, 135; *Howeana*, 141; *irregularis*, 65, 141; *Johnsoniana*, 141; *mammillata*, 129; *melanosperma*, 130, 131; *merdaria*, 138; *micropoda*, 141; *obdu-rata*, 131; *obturata*, 141; *paradoxa*, 129; *pseudocyanea*, 140; *rugomarginata*, 190; *rugoso-annulata*, 139; *Schraderi*, 141; *semiglobata*, 132-134; *siccipes*, 132; *siccipes radicata*, 132; *squamosa*, 126, 127; *stercoraria*, 133; *subbadia*, 139; *submerdaria*, 138; *tenuis*, 137; *umbilicata*, 142; *umbonatescens*, 128
- Strumella coryneoidea*, 199
- Studies in tropical ascomycetes — I, 235
- Substratum, A new lichen from an unusual, 95
- Syncephalastrum*, 147; *cinereum*, 149; *racemosum*, 149
- Syncephalis*, 147; *cordata*, 147; *nodosa*, 147; *pycnosperma*, 147; *Wynneae*, 147
- Synchytrium*, 151, 167; *decipiens*, 148, 155, 156, 158, 160, 163, 167; *endobioticum*, 149; *Puerariae*, 147, 156, 158; *pulverum*, 337; *Succisae*, 149; *Taraxaci*, 147
- Synedrella nodiflora*, 115
- Take-all of wheat, The *Ophiobolus* causing, 30
- Tanaka, T., New Japanese fungi. Notes and translations — XI, 81; —XII, 282
- Tetranthus hirsutus*, 116
- Thalictrum occidentale*, 177
- Thaxter, R., *Reliquiae Farlowianae*, 99
- Thlephora palmata*, 178
- Thelocarpon*, 95; *fimicola*, 96
- Thomas, H. E., Fitzpatrick, H. M., and Kirby, R. S., The *Ophiobolus* causing take-all of wheat, 30
- Thuja plicata*, 179
- Thyella*, 299
- Toronto Meeting, 50
- Trametes*, 183; *carnea*, 181, 183; *heteromorpha*, 184; *hispida*, 184; *sua-veolens*, 46, 47; *tenuis*, 6, 183
- Tremella frondosa*, 177; *lutescens*, 177; *mesenterica*, 177; *violacea*, 177
- Trichia*, 158
- Trichobasis Balsamorhizae*, 112; *Wyethiae*, 112
- Trichoglossum hirsutum*, 176
- Tricholoma*, 187; *bicolor*, 187; *per-sonatum*, 187; *subannulata*, 187
- Tridax procumbens*, 116
- Trinidad, Uredinales collected by Fred J. Seaver in, 12
- Trogia crispa*, 188
- Tropical ascomycetes — I, Studies in, 235
- Tsuga*, 181; *heterophylla*, 179
- Tubaria silvatica*, 269
- Tuber*, 176; *candidum*, 176
- Tuckahoe, Some characters of the southern, 222
- Tulostoma*, 50
- Tylopilus alboater*, 25
- Tylostoma*, 50; *campestre*, 50; *ver-rucosum*, 50, 232
- Tympanis*, 101
- Typhula phacorrhiza*, 178
- Ulmus*, 5
- Underwoodia*, 316
- Uredinales collected by Fred J. Seaver in Trinidad, 12
- Uredo abdita*, 120; *Balsamorhizae*, 112; *Ipomoeae*, 297; *Nyssae*, 336; *Parthenii*, 109; *ribicola*, 299; *Terebinthinaceae*, 251; *triannulata*, 111
- Urocystis Agropyri*, 279, 281
- Urocystis Agropyri* on redtop, 279
- Uromyces Alopecuri*, 229, 230; *Scirpi*, 229; *sediciosus*, 230; *triannula-tus*, 111
- Urophlyctis pluriannulatus*, 174
- Urticastrum*, 336
- Ustilaginoidea Oryzae*, 87; *Sacchari-Narengae*, 87
- Ustilago formosana*, 89; *Panici-pro-liferi*, 89
- Ustulina zonata*, 91
- Vaucheria*, 166, 167
- Venturia fimbriata*, 233
- Verbesina*, 113, 244, 245, 249, 250; *virginica*, 247-250, 298, 303, 304
- Vernonia*, 250; *angustifolia*, 304; *Blodgettii*, 248, 304; *flaccidifolia*, 248, 250, 303, 304; *glauca*, 248, 249, 298, 303-306; *noveboracensis*, 247, 248, 254, 303, 304, 306; *oligantha*, 248; *scariosa*, 104; *triflos-culosa*, 105
- Verticillium pyramidale*, 103
- Viguiera*, 244, 245; *deltioidea Parishii*, 110
- Wann, F. B., and Muenscher, W. C., A preliminary list of the myxomy-

- cetes of the Cayuga Lake Basin, 38
Wheat, The *Ophiobolus* causing take-all of, 30
Woronina, 146, 147, 151
Wyethia, 112
Ximenesia encelioides, 116
Xylaria, 311, 316; Longiana, 174
Zea Mais, 84
Zeller, S. M., Contributions to our knowledge of Oregon fungi—I, 173
Zinnia tenuiflora, 116
Zonanthemus, 115
Zygorhizidium Willei, 148, 151

I N D E X

UM